

# **Guidelines for use of G-CSF in Adult Haematology and Oncology Patients**

**Pathway of Care**

**Core Network Team**

Publication date	February 2021
Expected review date	February 2023
Version number	6
Version status	Final

## TABLE OF CONTENTS

---

1.0	INTRODUCTION.....	3
2.0	PROPHYLACTIC USE OF G-CSF IN CONJUNCTION WITH CHEMOTHERAPY GIVEN WITH CURATIVE INTENT .....	3
2.1	PRIMARY PROPHYLAXIS (I.E. USE WITH FIRST COURSE OF CHEMOTHERAPY ONWARDS).....	3
2.2	SECONDARY PROPHYLAXIS (I.E. USE AFTER EPISODE OF FN OR SEVERE NEUTROPENIA (GRADE 4) IN PRECEDING COURSE).....	4
3.0	PROPHYLACTIC USE OF G-CSF IN CONJUNCTION WITH PALLIATIVE CHEMOTHERAPY.....	5
4.0	THERAPEUTIC USE OF G-CSF.....	5
5.0	THE USE OF G-CSF IN SPECIFIC HAEMATOLOGICAL INDICATIONS.....	6
6.0	CHOICE AND DOSES OF G-CSF.....	6
7.0	DURATION AND TIMING OF THERAPY WITH G-CSF.....	7
8.0	IN CLINICAL TRIAL PATIENTS .....	7
9.0	APPROVED PRESCRIBER .....	7
10.0	APPENDIX A: DECISION MAKING ALGORITHM FOR PRIMARY GCSF PROPHYLAXIS.....	8
11.0	BIBLIOGRAPHY.....	9
12.0	PERSONNEL AND CONTACT INFORMATION.....	9
13.0	GLOSSARY .....	9
14.0	DOCUMENT ADMINISTRATION .....	10

## 1.0 INTRODUCTION

G-CSF guidelines have been developed in order to have a degree of consistency across the Kent and Medway Cancer Collaborative (K&MCC). In addition, there is a need to better define those patients who are most likely to benefit from G-CSF support. A reduction in febrile neutropenia (FN) is an important clinical outcome that justifies the use of G-CSF in the following indications. This guidance does not cover the use of G-CSF in clinical trials, or for the purpose of peripheral blood stem cell mobilization.

This guidance does not cover the use of G-CSF in children and young persons with cancer.

## 2.0 PROPHYLACTIC USE OF G-CSF IN CONJUNCTION WITH CHEMOTHERAPY GIVEN WITH CURATIVE INTENT

### 2.1 Primary Prophylaxis (i.e. use with first course of chemotherapy onwards)

- G-CSF is not recommended with most first line therapies.
- G-CSF should not be used to increase dose intensity outside of a clinical trial setting
- It is recommended that G-CSF primary prophylaxis is only given when chemotherapy is administered with curative intent (adjuvant / neo-adjuvant treatment). However, there may be instances where because of a high rate of febrile neutropenia (published or established through robust local audit) it is appropriate to give G-CSF as primary prophylaxis to patients receiving palliative chemotherapy.

**Table 1: Recommendations for G-CSF primary prophylaxis\***

<p>1. Where the incidence of FN associated with treatment is &gt; 20%. It should be noted that wherever possible, regimens of equal efficacy but with a lower risk of FN should be used. *</p>	<p>Examples of haematology regimens</p>	<ul style="list-style-type: none"> <li>• DHAP</li> <li>• ESHAP</li> <li>• Hyper C-VAD</li> <li>• CAIP</li> <li>• (R)-ICE</li> <li>• CODOX- M / IVAC</li> <li>• ICE</li> <li>• (R) CHOP 14</li> <li>• Minibeam</li> </ul>
	<p>Examples of oncology regimens</p>	<ul style="list-style-type: none"> <li>• FEC-T (breast)</li> <li>• FEC100</li> <li>• EC-Accelerated Paclitaxel</li> <li>• EC</li> <li>• TPF (H&amp;N)</li> <li>• <b>TP (H&amp;N)</b></li> <li>• Etoposide/ Platinum containing regimens (germ cell)</li> <li>• Etoposide/ platinum regimens (small cell cancers)</li> <li>• Docetaxel (2<sup>nd</sup> line NSCLC)</li> <li>• Nintedanib &amp; Docetaxel (NSCLC)</li> <li>• Folfirinox (upper GI)</li> </ul>

2. In patients with acute leukaemia in accordance with the guidelines below ( <a href="#">see section 4.0</a> )		
3. Primary prophylaxis may also be considered when used with regimens with febrile neutropenia rate of 10-20% in patients with one or more of the following risk factors; <ul style="list-style-type: none"> <li>● Extensive prior treatment</li> <li>● Older patients (&gt;65yrs)</li> <li>● The presence of open wounds or recent surgery</li> <li>● Advanced disease stage</li> <li>● Poor performance status where the use of chemotherapy is justified</li> <li>● Previous episodes of FN requiring admission</li> <li>● Cytopenias due to bone marrow involvement or pre-existing marrow <ul style="list-style-type: none"> <li>○ insufficiency (including previous irradiation to large volume of bone marrow)</li> </ul> </li> <li>● Poor nutritional status</li> <li>● Active infections including HIV</li> <li>● Serious co-morbidities</li> </ul>	Examples of haematology regimens	<ul style="list-style-type: none"> <li>● FCR (like regimens)</li> <li>● (R)-CHOP (like regimens)</li> <li>● Salvage regimens for lymphoma</li> </ul>
	Examples of oncology regimens	<ul style="list-style-type: none"> <li>● Docetaxel / Capecitabine</li> <li>● CAV</li> </ul>

\* N.B. While published rates of febrile neutropenia with platinum / etoposide combinations may not always exceed 20%, local experience has shown a high level of admission due to febrile neutropenia with these regimens. Local audit has shown rates of febrile neutropenia following docetaxel for 2<sup>nd</sup> line NSCLC exceed 20%.

Guidance during the COVID pandemic: The interim Covid NICE TA (<https://www.nice.org.uk/guidance/ng161>) has a lower threshold for primary prophylaxis (10% rather than usual 20% or 10-20% in select groups).

See [Appendix A](#) for a decision making algorithm for primary G-CSF prophylaxis

## 2.2 Secondary Prophylaxis (i.e. use after episode of FN or severe neutropenia (grade 4) in preceding course)

- Use of G-CSF to maintain dose intensity is not recommended in preference to dose reduction after prolonged neutropenia or an episode of neutropenic sepsis, except when chemotherapy is given with curative intent.

### 3.0 PROPHYLACTIC USE OF G-CSF IN CONJUNCTION WITH PALLIATIVE CHEMOTHERAPY

---

- G-CSF should not routinely be used in this indication ([see section 2.1](#))

### 4.0 THERAPEUTIC USE OF G-CSF

---

G-CSF must not be routinely prescribed for the treatment of patients with uncomplicated FN (duration of fever <10 days) or afebrile neutropenia.

G-CSF may be prescribed for the supportive treatment of patients with a high risk of infection-associated complications or severe FN (ANC <0.1 x 10<sup>9</sup>/l). G-CSF should continue until ANC >1.0 x 10<sup>9</sup>/l for 2 consecutive days and the expected neutrophil nadir has passed. High risk features include:

- Uncontrolled primary disease
- Pneumonia
- Clinically unwell with signs such as hypotension or organ dysfunction indicating potential risk of septic shock
- Expected prolonged duration of neutropenia (>10 days)
- Proven or suspected invasive fungal infection
- Age >65 years old
- Being hospitalised at the time of developing fever

<b>SOP No</b>	N/A	<b>Version</b>	6	<b>Supersedes version</b>	5	<b>Page 5 of 11</b>
<b>Written By</b>	Kate Miller	<b>Authorised by</b>	KMCC Chemo Group	<b>Date</b>	February 2021	
<b>KMCC document: No responsibility will be accepted for the accuracy of this information when used elsewhere.</b>						

## 5.0 THE USE OF G-CSF IN SPECIFIC HAEMATOLOGICAL INDICATIONS

**Table 2**

Indication	Recommendations	Initiation and duration of G-CSF
<b>AML</b>	G-CSF should not be used for priming of leukaemia cells (except with FLAG chemotherapy)	-
	G-CSF may be used following induction and for patients in remission following consolidation chemotherapy	Start when ANC < 0.5 until ANC >1.0 for 2 consecutive days NB Pegylated G-CSF should not be used in this group of patients
<b>MDS (high risk)</b>	G-CSF should only be considered for intermittent use in patients with severe neutropenia who experience recurrent infection Prolonged or continuous treatment with G-CSF is not recommended	Start when ANC <0.1. Stop when ANC >1.0 for 2 consecutive days
<b>MDS (low risk)</b>	G-CSF use in combination with epoetin should only be used following discussion within MDT	Three times a week
<b>ALL</b>	G-CSF should only be used where there is delayed recovery or infection following first few days of chemotherapy of initial induction or first post remission course of chemotherapy G-CSF should not be used for refractory or relapsed ALL	From day 5-8 post chemotherapy until ANC >1.0 for 2 consecutive days
<b>Aplastic anaemia/ inherited BM failure</b>	Use of G-CSF is not recommended	-

## 6.0 CHOICE AND DOSES OF G-CSF

- This guidance does not provide any recommendation on the formulation of G-CSF to be used.
- Pegylated G-CSF is considerably more expensive and should only be considered for patients unable to tolerate G-CSF. Please consult local formulary and pharmacy guidance.
- The usual dose of G-CSF in adult patients is 300 micrograms subcutaneously each day. A dose of 480 micrograms if the patient is >80kg.

## 7.0 DURATION AND TIMING OF THERAPY WITH G-CSF

---

- The data is too sparse to be absolutely specific about the most appropriate time to start G-CSF and the optimum duration of treatment. Reference should be made to the chemotherapy prescription within the cancer electronic prescribing system and the associated KMCC protocol if available.
- G-CSF should not usually be administered less than 24 hours following cytotoxic chemotherapy and is usually started 1-3 days after administration of myelotoxic chemotherapy. Administration of G-CSF daily for 5 days is usually adequate.

## 8.0 IN CLINICAL TRIAL PATIENTS

---

- Use of G-CSF should follow the trial protocol, irrespective of local policy.

## 9.0 APPROVED PRESCRIBER


---

- Oncology and haematology consultants and SpR's should initiate treatment. SHOs should only prescribe under the instruction of a consultant or SpR.

<b>SOP No</b>	N/A	<b>Version</b>	6	<b>Supersedes version</b>	5	<b>Page 7 of 11</b>
<b>Written By</b>	Kate Miller	<b>Authorised by</b>	KMCC Chemo Group	<b>Date</b>	February 2021	
<b>KMCC document: No responsibility will be accepted for the accuracy of this information when used elsewhere.</b>						

## 10.0 APPENDIX A: DECISION MAKING ALGORITHM FOR PRIMARY G-CSF PROPHYLAXIS

### Decision Making Algorithm for Primary G-CSF Prophylaxis



This decision making algorithm has been reproduced with kind permission from the London Cancer New Drugs Group.



## 11.0 BIBLIOGRAPHY

- ◆ Smith et al  
2006 Update of recommendations for the use of white blood cell growth factors: an evidence-based clinical practice guideline  
J Clin Oncol (2006) 24 (19) 3187-3205
- ◆ London Cancer New Drugs Group (January 2010)  
Guidance on the use of prophylactic granulocyte colony stimulating factor (GCSF) to support chemotherapy administration
- ◆ Information on file Amgen issued March 2010
- ◆ Smith et al 2015. Recommendations for the use of WBC Growth Factors: American Society of Clinical Oncology Clinical Practice Guideline Update.
- ◆ J Clin Oncol (2015) 33 (28) 3199-3212
- ◆ NHS England interim treatment options during the COVID-19 Pandemic (20/11/20)

## 12.0 PERSONNEL AND CONTACT INFORMATION

A comprehensive, up to date list of MDM contact details can be found on the KMCC website.

## 13.0 GLOSSARY

Acronyms in common usage throughout KMCC documentation

BNF	British National Formulary
BOPA	British Oncology Pharmacist Association
CNB	Cancer Network Board
COSHH	Control of substances hazardous to health regulations.
CYP	Children & Young People (in relation to the IOG)
DCCAG	Diagnostic Cross Cutting Advisory Group
DOG	Disease Orientated Group (NSSG/TSSG/TWG)
DVH	Darent Valley Hospital
DGT	Dartford and Gravesham NHS Trust
EK	East Kent
EKHUFT	East Kent Hospitals University Foundation Trust
EPS	Electronic Prescribing System
FP10(HNC)	Prescriptions issued by hospital doctors for dispensing in the community
GP	General Practitioner
HoP	High Level Operational Policy
IOSC	Improving Outcomes: A Strategy for Cancer
IV	Intravenous
K&C	Kent & Canterbury Hospital, Canterbury, (EKHUFT)
KMCC	Kent & Medway Cancer Collaborative
KMCRN	Kent & Medway Cancer Research Network
KOMS	Kent Oncology Management System

LSESN	London & South East Sarcoma Network
MFT	Medway Foundation Trust
MTW	Maidstone & Tunbridge Wells NHS Trust
NHS	National Health Service
NMP	Non-medical prescriber
NPSA	National Patient Safety agency
NOG	Non Surgical Oncology Group <i>(Permanent oncologist sub group of the DOGs with a specific responsibility for chemo/rad pathways and advice to the DOG, Network and GEOGRAPHICAL LOCATIONS on new drugs)</i>
PoC	Pathway of Care <i>(Network agreed disease site specific clinical guidelines)</i>
QEQM	Queen Elizabeth the Queen Mother Hospital, Margate (EKHUFT)
QoL	Quality of life
QSI	Quality service information system
QST	Quality Surveillance Team
RAT	Research and Trial Group <i>(Permanent sub-group of the DOGs with a specific responsibility for taking forward the clinical trials agenda)</i>
RMH	Royal Marsden Hospital
RNOH	Royal National Orthopaedic Hospital
SACT	Systemic Anti-Cancer therapy
SACT regimen	Systemic Anti-cancer prescription on the electronic prescribing system
SACT protocol	Systemic Anti-cancer protocol on KMCC website
TTO	Treatment to take home
QVH	Queen Victoria Foundation Trust Hospital East Grinstead
UCLH	University College Hospital London
WHH	William Harvey Hospital, Ashford (EKHUFT)
WK	West Kent

## 14.0 DOCUMENT ADMINISTRATION

<b>Document Title</b>	Oncological Treatment Guidelines for use of G-CSF in Adult Haematology and Oncology Patients
<b>Principal author</b>	Kate Miller
<b>Co-author(s)</b>	Caroline Waters / KMCC Chemo Group
<b>Current version number</b>	6
<b>Current status</b>	Final

The document is located in the Kent and Medway Cancer Network office, in hardcopy and in electronic format at [www.kmcc.nhs.uk/kent-and-medway-cancer-collaborative-kmcc/](http://www.kmcc.nhs.uk/kent-and-medway-cancer-collaborative-kmcc/)

<b>Enquiries:</b>	Caroline Waters. Lead Oncology Pharmacist, Kent and Medway Cancer Collaborative
<b>Date of Next Review:</b>	February 2023

### Revision History

Date of revision	New Version Number	Nature of Revision	Confirmation of Accuracy by
03/2006	001	Updated in accordance with ASCO 2006 guidelines	Wasifa Webb
20/03/08	002	Comments from HOG meeting of 24/04/08 incorporated	Drs Harper-Wynne and Rassam / Caroline Waters
15/05/08	003	Comments from HOG meeting of 29/05/08 incorporated	C Waters
29/05/08	004	Comments from Care Group relating to primary prophylaxis	C Waters
12/06/08	005		C Waters
12/06/08	1	Changes to section 1.0, table 1, section 5.0 (now section 6) and the addition of appendix 1	C Waters
01/03/10	1.1	Changes made to section 2.1 & 5.0 following HOG	C Waters
30/04/10	1.2	Changes made to section 3.10, 7.0 and appendix A following NCG	C Waters
June 2010	1.3	Final changes to section 4.0 following last consultation (e-mail)	C Waters
June 2010	1.4		C Waters
June 2010	2		C Waters
Oct- Dec 2012	2.1-2.3	Updated in accordance with ASCO 2006 guidelines	C Waters
January 2013	3		C Waters
May 2018	4	Updated in accordance with ASCO 2015 guidelines	K Miller
June 2018	4.1	Nintedanib & docetaxel and EC added to point 1 Table 1	C Waters
June 2018	5	Published following consultation with KMCC Chemotherapy Groups (circulated via email)	
Oct 2020	5.1	Update to section 6.0 Circulated to KMCC chemotherapy group for consultation via email by M Archer	Soleman Patel
Feb 2021	5.2	Update section 2.1 COVID interim dosing Section 5, Section 6, Section 7 Following email response	M.Archer
Feb 2021	5.2.1	All annotations removed ready for reformatting. correction of abbreviations to KMCC from KMCN within document Added up to date glossary Reformatted by R Patel	M.Archer
Feb 2021	6	Published following consultation with KMCC Chemotherapy Groups	M.Archer