

| | |
|--|--|
| Indication | Glioma |
| Treatment Intent | Palliative |
| Frequency and number of cycles | Repeat every 42 days for the maximum of 6 cycles. |
| Monitoring Parameters pre-treatment | <ul style="list-style-type: none"> • Monitor LFT's, U&E's Glucose and FBC at each cycle. • If Plts ≥ 150 and neuts ≥ 1.5 proceed at full dose. • If neuts ≥ 1.5 and Plts 100-149 discuss with consultant. • If neuts < 1.5 and/or PLT < 100 defer chemo. • Repeat FBC weekly. If FBC still low after first delay, discuss with consultant and consider dose reduction. • Dose reduction should be considered if grade 3 or 4 non-haematological toxicity or repeat appearance of grade 2 (except N&V and alopecia). Delay until resolution of toxicity to \leq grade 1. • Lung function-as clinically indicated. • *Dose can be increased to $120\text{mg}/\text{m}^2$ at consultant's discretion. • Hepatic impairment:– d/w consultant • Renal impairment: <ul style="list-style-type: none"> ○ If CrCl $> 60\text{ml}/\text{min}$ give 100% dose ○ $45\text{-}60\text{ml}/\text{min}$ reduce to 75% ○ $30\text{-}44\text{ml}/\text{min}$ reduce to 50%. ○ Not recommended if CrCl $< 30\text{ml}/\text{min}$. |
| References | KMCC proforma BRA-006v1, The North London Cancer Network Dosage Adjustment for Cytotoxics in Hepatic Impairment 2009, The North London Cancer Network Dosage Adjustment for Cytotoxics in Renal Impairment 2009 SPC accessed online 05/11/2019 |

NB For funding information, refer to CDF and NICE Drugs Funding List

Repeat every 42 days

| Day | Drug | Dose | Route | Administration |
|-----|-------------------------|---|-------|---|
| 1 | LOMUSTINE (CCNU) | $100\text{mg}/\text{m}^2$ (see notes above *) | PO | Take as a single dose at night on Day 1 ONLY. Available as 40mg capsule |
| | Ondansetron | 8mg | PO | Take BD for 3 days. Take the first dose 30 minutes before taking the Lomustine capsules. |
| | Domperidone | 10mg | PO | Up to TDS PRN. Maximum 30mg day. Do not take for more than 7 days continuously. |
| | Dexamethasone | 6mg | PO | OM for 3/7 |
| | Movicol sachet | 1 sachet | PO | Take the contents of ONE sachet dissolved or mixed with water BD as required. |

| | | | |
|--------------------|----------|---|---------------------|
| Protocol No | BRA-006 | Kent and Medway SACT Protocol Disclaimer: No responsibility will be accepted for the accuracy of this information when used elsewhere. | |
| Version | V2 | Written by | M.Archer |
| Supersedes version | V1 | Checked by | C.Waters E.Parry |
| Date | 05/12/19 | Authorising consultant (usually NOG Chair) | J.Glendenning |