

Indication	ALL
Treatment Intent	Palliative for the elderly
Frequency and number of cycles	35 days Maximum of 2 cycles
Monitoring Parameters pre-treatment	<ul style="list-style-type: none"> FBC, U&Es and LFTS before each cycle. Haematological - Give first cycle regardless of FBC. Proceed with subsequent cycles when neutrophils > 1.0 x 10⁹/L and platelets > 50 x 10⁹/L. <u>Dose modification:</u> <u>Vincristine:</u> Bilirubin 26-51µmol/L or AST/ALT 60-180 units give 50%; bilirubin > 51µmol/L and AST/ALT normal give 50%; bilirubin > 51µmol/L and AST/ALT > 180 units omit. Neurotoxicity - Grade 2 motor and Grade 3 sensory toxicity give Vincristine 50% dose. Antifungal cover has not been included in this regime due to interaction with of fluconazole and vincristine. If patients are high risk Consultants can consider Liposomal amphotericin prophylaxis instead.
References	UKALL 10 protocol and UKALL 12 protocol HOG minutes 07/03/19

NB For funding information, refer to CDF and NICE Drugs Funding List

Repeat every 35 days

Day	Drug	Dose	Route	Infusion Duration	Administration
1, 8, 15 and 22	Vincristine	1.4mg/m² Max dose 2mg	IV	5-10 minutes	In 50ml Sodium Chloride 0.9% FOR INTRAVENOUS USE ONLY. FATAL IF GIVEN BY ANY OTHER ROUTE

Protocol No	HAEM-ALL-013	Kent and Medway SACT Protocol Disclaimer: No responsibility will be accepted for the accuracy of this information when used elsewhere.	
Version	V1	Written by	M.Archer
Supersedes version	New protocol	Checked by	C.Waters K.Miller
Date	02/07/20	Authorising consultant (usually NOG Chair)	K.Yip

Dispense on day 1 of cycle:

TTO	Drug	Dose	Route	Directions
	Prednisolone	40mg/m ²	PO	Daily for 28 days and then reduce as directed over 7 days. Take in 2 divided doses. Take with or after food.
	Metoclopramide	10mg	PO	TDS when required. Do not take for more than 5 days continuously.
	Allopurinol	300mg Dose reduce in renal impairment.	PO	OD Take with or after food with a full glass of water. Cycle 1 only.
	Co-trimoxazole	480mg	PO	BD on a Monday, Wednesday and Friday only.
	Omeprazole	20mg	PO	OD
	Chlorhexidine mouthwash	10ml	mouthwash	QDS

Protocol No	HAEM-ALL-013	Kent and Medway SACT Protocol Disclaimer: No responsibility will be accepted for the accuracy of this information when used elsewhere.	
Version	V1	Written by	M.Archer
Supersedes version	New protocol	Checked by	C.Waters K.Miller
Date	02/07/20	Authorising consultant (usually NOG Chair)	K.Yip